



# Town of Darien

David Knauf, MPH, MS, REHS

Director of Health

Phone: (203) 656-7320

e-mail: [dknauf@darienct.gov](mailto:dknauf@darienct.gov)

[www.darienhealth.com](http://www.darienhealth.com)

## Health Department

### It's That Time of Year!

### TICKS ARE COMING OUT IN LARGE NUMBERS

The black legged or “deer” tick (*Ixodes scapularis*) is well known for spreading Lyme disease. Now, after a long winter, the ticks are out in large numbers, looking for a blood meal, possibly from you! So here are some tips and things to know to protect yourself and your family:

#### TICK FACTS

- Ticks are external bloodsucking parasites that feed on humans or other mammals. They are entirely dependent on blood or tissue fluids for life.
- Ticks are second only to mosquitoes as transmitters of infectious agents to humans. The longer an infective tick is permitted to feed, the greater chances of transmitting the infection.
- A tick seeks a blood meal by climbing onto vegetation and using its forelegs to feel and grab onto a host. They are found from ground level to about three feet above. Carbon dioxide, scent, body heat and other stimuli attract ticks to a host.
- The female requires a blood meal in order to lay its eggs.
- Male ticks do not transmit Lyme Disease.
- Many tick life cycles are completed within one or two years depending upon the ease or difficulty in finding a blood meal.

#### PERSONAL PROTECTION

These techniques will reduce your chance of a tick bite:

- Avoid tick-infested areas wherever possible. Stay on established paths and try not to brush against bushes where ticks may be waiting for you and their next meal.
- Avoid sitting on the ground.
- Wear light colored clothing. This makes it easier to spot ticks on your clothing.
- Wear long-sleeved shirt and long trousers. This will reduce the skin area exposed to ticks.
- Tuck your shirt into your trousers and your trouser legs into your socks. This keeps the ticks on the outside of your clothing as they crawl upward.
- Use an EPA-approved tick repellent and wash it off when you return home.
- Treat all the shoes your family might wear in tick habitat on the first of every month, May - August with permethrin clothing repellent spray.
- Conduct frequent tick checks. This includes a visual inspection of the clothing and exposed skin, followed by a naked, full body examination in private. Look and feel all over your skin and scalp for new bumps.
- Take a shower and wash your hair when you come inside.
- Ticks can get on your pets so check them regularly, especially around the ears and eyes.

## PROPERTY MANAGEMENT

Taking the following actions will reduce but not eliminate the number of ticks that may be on your property:

- Ticks are susceptible to dehydration. You can reduce humidity by: pruning trees, clearing brush, removing litter, mowing grass and allowing grass to dry between watering. Also, move the stubby overgrowth between the lawn and the woods farther back from areas frequented by people and pets.
- Modify your property so it is less attractive to animals that are hosts to ticks by: eliminating bird feeders and baths, salt licks (for deer), clear away wood and brush piles, garbage, leaf piles and removing stone walls.
- Directly kill ticks that live on the property by applying acaricides (tick-killing pesticides). Follow the directions carefully as these are toxic. Better still, use an exterminator.
- Modify outdoor areas: widen trails and move play areas to safer, more open places.

## TICK REMOVAL

***Free tick removal kits are available at the Darien Health Department!***

- Using fine pointed tweezers, grasp the tick's head as close to the patient's skin as is possible and apply steady, gentle upward pressure until the tick releases its hold. Do not yank! Part of the head may remain embedded in the skin. Put the tick into a small plastic sandwich type bag and bring it to the Darien Health Dept. for testing. Wash the bite site and the tweezers with suitable antiseptic. Notify your physician.
- Children should be told to seek adult help for tick removal. It is safer to wait and remove it properly than to pull it off with your fingers. Mouthparts may remain embedded in the skin.
- Do not prick, crush or burn the tick, as it may cause the release of infected tick fluid.
- Do not try to smother the tick (e.g. petroleum jelly, nail polish) as the tick has enough oxygen to complete the feeding and this tactic could make removing the tick even more difficult.

## LYME DISEASE SYMPTOMS

After the removal of the tick, submit it to the Health Department for testing & watch for any of the following:

- Flu-like feeling, aches, fever, headache.
- Enlarging rash at the original site of the bite; reddish on light skin, bruised looking on darker skin. About 50 percent of light skinned adults get a rash and some get multiple rashes. Joint swelling or pain may also occur.

If any of the above symptoms occur, or if the tick test comes back positive, notify your physician immediately. Also, you must be aware of the fact that symptoms of Lyme Disease can vary between patients so physicians must be aware of your activities and potential exposure to ticks.

For more information, call the **Darien Health Department at (203) 656-7320**, visit the website at [www.darienhealth.com](http://www.darienhealth.com) or drop in at the **Town Hall, Room 109**.